

CLASSIFICATION R. TRI 200

CENTRAL INTELLIGENCE AGENCY

**RESTRICTED**  
**INFORMATION REPORT**

## REPORT

STAT

COUNTRY : USSR

DATE DISTR. 13 September 1948

**SUBJECT**      Scientific Research

NO. OF PAGES

PLACE ACQUIRED USSR

NO. OF ENCLS.  
(LISTED BELOW)

DATE OF  
INFORMATION 1946-47

SUPPLEMENT TO  
REPORT NO.

STAT

THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE OF THE UNITED STATES WITHIN THE MEANING OF THE ESPIONAGE ACT 50 U.S.C. 31 AND 32, AS AMENDED. ITS TRANSMISSION OR THE REVELATION OF ITS CONTENTS IN ANY MANNER TO AN UNAUTHORIZED PERSON IS PROHIBITED BY LAW. REPRODUCTION OF THIS FORM IS PROHIBITED. HOWEVER, INFORMATION CONTAINED IN FOOT OF THE FORM MAY BE UTILIZED AS DEEMED NECESSARY BY THE RECEIVING AGENCY.

THIS IS UNEVALUATED INFORMATION FOR THE RESEARCH  
USE OF TRAINED INTELLIGENCE ANALYSTS

**SOURCE** Documentary as indicated. (Information specifically requested.)

RECENTLY PUBLISHED RESEARCH OF THE  
DNEPROPETROVSK METALLURGICAL INSTITUTE, USSR

"Kinetics of the Decomposition of Tetrafluoroborates in Aqueous Solutions," I. G. Ryss, M. M. Slutskaya, Lab Gen Chem, Dnepropetrovsk Metal Inst imeni Stalin, 12 pp

"Zhur Fiz Khim" Vol XXI, no 5, May 1947

Discusses, with detailed tables, illustrations and formulas, the kinetics of decomposition with result that the constant of the rate  $k' = 0.4343K$  at temperature of 20 to 80°C. Rate of reaction was found to increase with rise in temperature. (181100)

"Crystallization Equilibrium of Solutions of Stannous Chloride," I. G. Ryss, E. Ya. Turkanov, Dnepropetrovsk Metal Inst

"Zhur Priklad Knim" Vol 19, 1947, pp 958-65.

In the system  $\text{SnCl}_2\text{-H}_2\text{O-HCl}$ , equilibrium is attained from both sides in 2-3 hours at  $50^\circ$ . From 0 to  $250^\circ$ , with up to 10% HCl, the solid phase is  $\text{SnCl}_2\cdot 2\text{H}_2\text{O}$ . Isotherms at those temperatures, with %  $\text{SnCl}_2\cdot 2\text{H}_2\text{O}$  i. solution plotted against % HCl, show a minimum around 5.5-8% HCl, the latter the higher the temperature. Cryometric determinations without HCl (other than that due to hydrolysis of  $\text{SnCl}_2$ ) gave: from a 34.7%  $\text{SnCl}_2$  solution, the crystallization

6. 1 -

**CLASSIFICATION**

RESTRICTED

CLASSIFICATION									
STATE	NAVY	NARB	DISTRIBUTION						
ARMY	AIR	EDB							

**RESTRICTED**

RESTRICTED

RESTRICTED

STAT

temperature of ice is  $-6.8^{\circ}$ ;  $\text{SnCl}_2$  content of the solution in equilibrium with the eutectic, 37.9%. From 36.8 and 37.3%  $\text{SnCl}_2$ , ice crystallizes at  $-6.25^{\circ}$  and  $-6.45^{\circ}$ , respectively; eutectic arrest at  $-6.8^{\circ}$ . Slightly moist  $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$  showed on heating a temperature arrest at  $41.4^{\circ}$ ; it could not be decided whether this corresponds to complete melting, nor could a second eutectic  $\text{SnCl}_2 \cdot 2\text{H}_2\text{O} - \text{SnCl}_2$  be established; it appears that  $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$  melts incongruently but with a composition very close to it.

"Hydrolytic Equilibria in Solutions of Sodium Fluosilicate at  $11^{\circ}$ ," I. G. Ryss, Dnepropetrovsk Metal Inst

"Zhur Obshch Khim" Vol 16, 1946, pp 331-40

The degree of hydrolysis of  $\text{Na}_2\text{SiF}_6$  at  $11^{\circ}$  at concentrations 0.02 M to 0.00048 M was determined. The following equilibrium constants were calculated:  
 $\text{SiF}_6^{2-} \rightleftharpoons \text{SiF}_4 + 2\text{F}^-$ ,  $K_1 = 0.65 \times 10^{-6}$ ;  
 $\text{SiF}_4 + 2\text{H}_2\text{O} \rightleftharpoons \text{SiO}_2 + 4\text{HF}$ ,  $K_2 = 1.04 \times 10^{-8}$ ;  $\text{SiF}_6^{2-} + 2\text{H}_2\text{O} \rightleftharpoons \text{SiO}_2 + 4\text{H}^+ + 6\text{F}^-$ ,  $K_3 = 5.4 \times 10^{-27}$ .  
 Generally speaking, the solutions of the hexafluoride reach hydrolytic equilibrium rapidly and further changes occur only very slowly.

- END -

- 2 -

RESTRICTED

RESTRICTED